

## A NOTE ON MAXIMAL NON-PRIME IDEALS

S. VISWESWARAN\* AND ANIRUDHDHA PARMAR

ABSTRACT. The rings considered in this article are commutative with identity  $1 \neq 0$ . We say that a proper ideal  $I$  of a ring  $R$  is a maximal non-prime ideal if  $I$  is not a prime ideal of  $R$  but any proper ideal  $A$  of  $R$  with  $I \subseteq A$  and  $I \neq A$  is a prime ideal. That is, among all the proper ideals of  $R$ ,  $I$  is maximal with respect to the property of being not a prime ideal. The concept of maximal non-maximal ideal and maximal non-primary ideal of a ring can be similarly defined. The aim of this article is to characterize ideals  $I$  of a ring  $R$  such that  $I$  is a maximal non-prime (respectively, a maximal non-maximal, a maximal non-primary) ideal of  $R$ .

### 1. INTRODUCTION

The rings considered in this article are nonzero commutative with identity. If  $R$  is a subring of a ring  $T$  with identity 1, then we assume that  $1 \in R$ . If a set  $A$  is a subset of a set  $B$  and  $A \neq B$ , we denote it symbolically using the notation  $A \subset B$ . Let  $P$  be a property of rings. Let  $R$  be a subring of a ring  $T$ . Recall from [4] that  $R$  is a maximal non- $P$ , if  $R$  does not have  $P$ , whereas each subring  $S$  of  $T$  with  $R \subset S$  has property  $P$ . The concept of maximal non-Noetherian subring of a ring  $T$  was investigated in [3]. There are other interesting research articles which appeared in the literature focussing on maximal non- $P$  subring of a ring  $T$  (see for example, [2, 4]). Let  $R$  be a non-zero commutative ring with identity. A proper ideal  $I$  of a ring  $R$  is said to be a maximal non-prime ideal of  $R$  if the following conditions hold: (i)  $I$  is not a

---

MSC(2010): Primary 13A15; Secondary 13C05.

Keywords: Maximal non-prime ideal, maximal non-maximal ideal, maximal non-primary ideal, maximal non-irreducible ideal.

Received: 12 June 2015, Accepted: 15 July 2015.

\*Corresponding author .

prime ideal of  $R$  and (ii) If  $A$  is any proper ideal of  $R$  such that  $A$  contains  $I$  properly, then  $A$  is a prime ideal of  $R$ . Similarly, we can define the concept of a maximal non-maximal (respectively, a maximal non-primary) ideal of  $R$ . Motivated by the above mentioned works on maximal non- $P$  subrings, in this article, we focus our attempt on characterizing maximal non-prime (respectively, maximal non-maximal, maximal non-primary) ideals of a ring  $R$ . Let  $I$  be a proper radical ideal of a ring  $R$ . It is proved in Proposition 3.2 that  $I$  is a maximal non-primary ideal of  $R$  if and only if  $I$  is a maximal non-prime ideal of  $R$  if and only if  $I$  is a maximal non-maximal ideal of  $R$  if and only if  $I = M_1 \cap M_2$  for some distinct maximal ideals  $M_1, M_2$  of  $R$ . Let  $I$  be a proper ideal of  $R$  such that  $I \neq \sqrt{I}$ . It is shown in Proposition 4.1 that  $I$  is a maximal non-prime ideal of  $R$  if and only if  $I$  is a maximal non-maximal ideal of  $R$  if and only if  $\sqrt{I} = M$  is a maximal ideal of  $R$  with  $M^2 \subseteq I$ , and  $M = Rx + I$  for any  $x \in M \setminus I$ . Moreover, it is proved in Proposition 4.2 that  $I$  is a maximal non-primary ideal of  $R$  if and only if  $\sqrt{I} = P$  is a prime ideal of  $R$  such that  $R/I$  is a quasilocal one-dimensional ring and  $P/I$  is a minimal ideal of  $R/I$ .

By a quasilocal ring we mean a ring which admits only one maximal ideal. A Noetherian quasilocal ring is referred to as a local ring.. By dimension of a ring  $R$ , we mean its Krull dimension and we use the abbreviation  $\dim R$  to denote the dimension of a ring  $R$ . We denote the nilradical of a ring  $R$  by  $\text{nil}(R)$ . A ring  $R$  is said to be reduced if  $\text{nil}(R) = (0)$ .

## 2. SOME PRELIMINARY RESULTS

As mentioned in the introduction the rings considered in this article are commutative with identity  $1 \neq 0$ . We begin with the following lemma.

**Lemma 2.1.** *Let  $R$  be a ring. If  $P_1, P_2$  are incomparable prime ideals of  $R$  under inclusion, then  $P_1 \cap P_2$  is not a primary ideal of  $R$ .*

*Proof.* Let  $I = P_1 \cap P_2$ . Since  $P_1$  and  $P_2$  are incomparable under inclusion, there exist  $a \in P_1 \setminus P_2$  and  $b \in P_2 \setminus P_1$ . Note that  $ab \in I$ . By the choice of  $a, b$ , it is clear that  $a \notin I$  and no power of  $b \in I$ . This proves that  $I = P_1 \cap P_2$  is not a primary ideal of  $R$ .  $\square$

**Lemma 2.2.** *Let  $R$  be a reduced ring which is not an integral domain. If every nonzero proper ideal of  $R$  is primary, then  $R$  has exactly two prime ideals and both of them are maximal ideals of  $R$ .*

*Proof.* Since  $R$  is reduced but not an integral domain, it follows that  $R$  has at least two minimal prime ideals. Let  $P_1, P_2$  be distinct minimal prime ideals of  $R$ . Now we obtain from Lemma 2.1 and the hypothesis that  $P_1 \cap P_2 = (0)$ . We prove that  $P_1, P_2$  are maximal ideals of  $R$ . Let  $M$  be a maximal ideal of  $R$  such that  $P_1 \subseteq M$ . We claim that  $M = P_1$ . Suppose that  $P_1 \neq M$ . Then  $M \not\subseteq P_1 \cup P_2$ . Let  $a \in M \setminus (P_1 \cup P_2)$  and  $b \in P_2 \setminus P_1$ . As  $ab \notin P_1$ , it follows that  $ab \neq 0$ . Hence  $Rab$  is a primary ideal of  $R$ . Note that  $Rab \subseteq P_2$ . Hence it follows from the choice of  $a$  that no power of  $a \in Rab$ . Therefore,  $b \in Rab$ . This implies that  $b = rab$  for some  $r \in R$  and so  $b(1 - ra) = 0$ . As  $b \notin P_1$ , it follows that  $1 - ra \in P_1 \subset M$ . From  $a \in M$ , we obtain that  $1 = 1 - ra + ra \in M$ . This is a contradiction. Therefore,  $P_1 = M$  is a maximal ideal of  $R$ . Similarly, it follows that  $P_2$  is a maximal ideal of  $R$ . From  $P_1 \cap P_2 = (0)$ , we get that  $R$  has exactly two prime ideals which are  $P_1$  and  $P_2$  and moreover, both are maximal ideals of  $R$ .  $\square$

**Lemma 2.3.** *Let  $R$  be a ring such that every nonzero proper ideal of  $R$  is primary. Then  $\dim R \leq 1$ . Moreover, if  $R$  is not a reduced ring, then  $R$  is necessarily quasilocal.*

*Proof.* Suppose that  $\dim R > 1$ . Then there exists a chain of prime ideals  $P_1 \subset P_2 \subset P_3$  of  $R$ . Let  $a \in P_2 \setminus P_1$  and  $b \in P_3 \setminus P_2$ . Since  $ab \notin P_1$ , it is clear that  $ab \neq 0$  and hence  $Rab \neq (0)$ . Observe that  $Rab \subseteq P_2$ . By hypothesis,  $Rab$  is a primary ideal of  $R$ . From the choice of the element  $b$ , it is clear that no power of  $b$  can belong to  $Rab$ . Hence  $a \in Rab$ . This implies that  $a = rab$  for some  $r \in R$  and so  $a(1 - rb) = 0$ . Since  $a \notin P_1$ , it follows that  $1 - rb \in P_1 \subset P_3$ . From  $b \in P_3$ , we obtain that  $1 = 1 - rb + rb \in P_3$ . This is a contradiction. Therefore,  $\dim R \leq 1$ .

We next prove the moreover assertion. Suppose that  $R$  is not quasilocal. Then there exist at least two distinct maximal ideals  $M_1, M_2$  of  $R$ . As we are assuming that  $R$  is not a reduced ring, it follows that  $M_1 \cap M_2 \neq (0)$ . Hence by hypothesis,  $M_1 \cap M_2$  is a primary ideal of  $R$ . This contradicts Lemma 2.1. Therefore,  $R$  is necessarily quasilocal.  $\square$

**Lemma 2.4.** *Let  $R$  be a ring which is not reduced. Suppose that  $(0)$  is not a primary ideal of  $R$ . If every nonzero proper ideal of  $R$  is primary, then  $\text{nil}(R)$  is a minimal prime ideal of  $R$ . Indeed,  $\text{nil}(R)$  is a minimal ideal of  $R$ .*

*Proof.* We know from Lemma 2.3 that  $R$  is necessarily quasilocal. Let  $M$  be the unique maximal ideal of  $R$ . Since  $R$  is not reduced,  $\text{nil}(R) \neq (0)$ . Hence  $\text{nil}(R)$  is a primary ideal of  $R$  and so it follows from [1,

Proposition 4.1] that  $\sqrt{\text{nil}(R)} = \text{nil}(R)$  is a prime ideal of  $R$ . Since  $\text{nil}(R) \subseteq P$  for any prime ideal of  $R$ , it follows that  $\text{nil}(R)$  is a minimal prime ideal of  $R$ . As  $(0)$  is not a primary ideal of  $R$ , it follows from [1, Proposition 4.2] that  $\sqrt{(0)}$  is not a maximal ideal of  $R$ . Thus  $\text{nil}(R) \subset M$ . We prove that for any nonzero  $a \in \text{nil}(R)$ ,  $\text{nil}(R) = Ra$ . First we verify that for any  $b \in \text{nil}(R) \setminus (0)$  and for any  $m \in M \setminus \text{nil}(R)$ ,  $bm = 0$ . Suppose that  $bm \neq 0$ . By hypothesis,  $Rbm$  is a primary ideal of  $R$ . In fact  $Rbm$  is a  $\text{nil}(R)$ -primary ideal of  $R$ . Since no power of  $m \in \text{nil}(R)$ , we obtain that  $b \in Rbm$ . This implies that  $b = rbm$  for some  $r \in R$ . Thus  $b(1 - rm) = 0$ . As  $1 - rm$  is a unit in  $R$ , it follows that  $b = 0$ . This is a contradiction. Hence for any nonzero  $b \in \text{nil}(R)$  and  $m \in M \setminus \text{nil}(R)$ ,  $bm = 0$ . Let  $x \in \text{nil}(R)$ . We assert that  $x \in Ra$ . This is clear if  $x = 0$ . If  $x \neq 0$ , then  $xm = 0 \in Ra$ . Now  $Ra$  is a  $\text{nil}(R)$ -primary ideal of  $R$  and no power of  $m \in \text{nil}(R)$ . Hence it follows that  $x \in Ra$ . This proves that for any nonzero  $a \in \text{nil}(R)$ ,  $\text{nil}(R) = Ra$ . This shows that  $\text{nil}(R)$  is a minimal ideal of  $R$ .  $\square$

**Lemma 2.5.** *Let  $R$  be a quasilocal ring with  $M$  as its unique maximal ideal. Suppose that  $R$  is not reduced and  $\text{nil}(R)$  is a prime ideal of  $R$  with  $\text{nil}(R) \neq M$ . If  $\text{nil}(R)$  is a minimal ideal of  $R$ , then  $(0)$  is not a primary ideal of  $R$ .*

*Proof.* Let  $a \in \text{nil}(R)$ ,  $a \neq 0$ . Let  $b \in M \setminus \text{nil}(R)$ . Since  $\text{nil}(R)$  is a simple  $R$ -module, it follows that  $M(\text{nil}(R)) = (0)$  and so  $ab = 0$ . Now  $a \neq 0$  and as  $b \notin \text{nil}(R)$ , it follows that  $b^n \neq 0$  for all  $n \geq 1$ . This proves that  $(0)$  is not a primary ideal of  $R$ .  $\square$

**Lemma 2.6.** *Let  $R$  be a ring which is not reduced. If every nonzero proper ideal of  $R$  is a prime ideal of  $R$ , then  $R$  is quasilocal with  $\text{nil}(R)$  as its unique maximal ideal and  $(\text{nil}(R))^2 = (0)$ . Moreover, for any  $x \in \text{nil}(R) \setminus \{0\}$ ,  $\text{nil}(R) = Rx$ .*

*Proof.* Since any prime ideal is primary, it follows from Lemma 2.3 that  $R$  is necessarily quasilocal. Let  $M$  be the unique maximal ideal of  $R$ . We prove that  $M = \text{nil}(R)$ . Let  $m \in M$ . We assert that  $m^2 = 0$ . Suppose that  $m^2 \neq 0$ . Then  $Rm^2$  is a prime ideal of  $R$ . Therefore,  $m \in Rm^2$ . This implies that  $m = rm^2$  for some  $r \in R$  and so  $m(1 - rm) = 0$ . From  $1 - rm$  is a unit in  $R$ , it follows that  $m = 0$ . This is a contradiction. Thus for any  $m \in M$ ,  $m^2 = 0$  and so  $M = \text{nil}(R)$ . Hence  $M$  is the only prime ideal of  $R$ . Let  $a, b \in M$ . We show that  $ab = 0$ . This is clear if either  $a = 0$  or  $b = 0$ . Suppose that  $a \neq 0$  and  $b \neq 0$ . Then  $Ra, Rb$  are prime ideals of  $R$ . Therefore,  $Ra = Rb = M$ . This implies that  $a = ub$  for some unit  $u \in R$ . It follows from  $b^2 = 0$  that  $ab = 0$ . This proves that  $M^2 = (\text{nil}(R))^2 = (0)$ .

We next prove the moreover part. Let  $x \in \text{nil}(R) \setminus \{0\}$ . Then  $Rx$  is a prime ideal of  $R$ . From the fact that  $\text{nil}(R)$  is the only prime ideal of  $R$ , it follows that  $\text{nil}(R) = Rx$ .  $\square$

### 3. RADICAL NON-MAXIMAL PRIME IDEALS

The aim of this section is to determine proper radical ideals  $I$  of a ring  $R$  such that  $I$  is a maximal non-prime ideal. We start with the following lemma.

**Lemma 3.1.** *Let  $D$  be an integral domain which is not a field. Then it admits nonzero proper ideals which are not prime ideals.*

*Proof.* Let  $d \in D$  be a nonzero nonunit. Then for any  $n \geq 2$ ,  $Dd^n$  is a proper nonzero ideal of  $D$  which is not a prime ideal of  $D$ .  $\square$

**Proposition 3.2.** *Let  $R$  be a ring and  $I$  be a proper radical ideal of  $R$ . Then the following statements are equivalent:*

- (i)  $I$  is a maximal non-primary ideal of  $R$ .
- (ii)  $I = M_1 \cap M_2$  for some distinct maximal ideals  $M_1, M_2$  of  $R$ .
- (iii)  $I$  is a maximal non-maximal ideal of  $R$ .
- (iv)  $I$  is a maximal non-prime ideal of  $R$ .

*Proof.* (i)  $\Rightarrow$  (ii) Note that  $R/I$  is a reduced ring and as  $I$  is not primary, it follows that  $I$  is not a prime ideal of  $R$  and so  $R/I$  is not an integral domain. Since  $I$  is a maximal non-primary ideal of  $R$ , it follows that every nonzero proper ideal of  $R/I$  is primary. Hence we obtain from Lemma 2.2 that there exist distinct maximal ideals  $M_1, M_2$  of  $R$  such that  $I = M_1 \cap M_2$ .

(ii)  $\Rightarrow$  (iii) We know from Lemma 2.1 that  $I = M_1 \cap M_2$  is not a primary ideal and hence it is not a maximal ideal of  $R$ . Let  $A$  be any proper ideal of  $R$  such that  $M_1 \cap M_2 \subset A$ . Then either  $A \not\subseteq M_1$  or  $A \not\subseteq M_2$ . Without loss of generality we may assume that  $A \not\subseteq M_1$ . Then  $A + M_1 = R$ . Hence  $1 = a + x$  for some  $a \in A$  and  $x \in M_1$ . Now for any  $y \in M_2$ ,  $y = ay + xy \in A + M_1M_2 = A$ . This proves that  $M_2 \subseteq A$  and so  $A = M_2$ . Thus the only proper ideals  $A$  of  $R$  which contain  $I$  properly are  $M_1$  and  $M_2$  and both are maximal ideals of  $R$ . Therefore, we obtain that  $I$  is a maximal non-maximal ideal of  $R$ .

(iii)  $\Rightarrow$  (iv) Let  $A$  be a proper ideal of  $R$  with  $I \subset A$ . Then by (iii)  $A$  is a maximal ideal of  $R$ . Hence  $A$  is a prime ideal of  $R$ . We claim that  $I$  is not a prime ideal of  $R$ . Suppose that  $I$  is a prime ideal of  $R$ . Since  $R/I$  is not a field, it follows from Lemma 3.1 that  $R/I$  admits nonzero proper ideals which are not maximal ideals. This contradicts (iii). Therefore,  $I$  is not a prime ideal of  $R$ . This shows that  $I$  is a maximal non-prime ideal of  $R$ .

(iv)  $\Rightarrow$  (i) Let  $A$  be any proper ideal of  $R$  with  $I \subset A$ . Then by (iv)  $A$  is a prime ideal and hence is a primary ideal of  $R$ . Since  $I$  is a radical ideal of  $R$  and is not a prime ideal of  $R$ , we get that  $I$  is not a primary ideal of  $R$ . This proves that  $I$  is a maximal non-primary ideal of  $R$ .  $\square$

#### 4. NON-RADICAL MAXIMAL NON-PRIME IDEALS

The aim of this section is to determine ideals  $I$  of a ring  $R$  such that  $I \neq \sqrt{I}$  and  $I$  is a maximal non-prime ideal of  $R$ .

**Proposition 4.1.** *Let  $I$  be a proper ideal of a ring  $R$  such that  $I \neq \sqrt{I}$ . Then the following statements are equivalent:*

- (i)  $I$  is a maximal non-prime ideal of  $R$ .
- (ii)  $\sqrt{I}$  is a maximal ideal of  $R$ ,  $(\sqrt{I})^2 \subseteq I$ , and  $\sqrt{I} = Rx + I$  for any  $x \in \sqrt{I} \setminus I$ .
- (iii)  $I$  is a maximal non-maximal ideal of  $R$ .

*Proof.* (i)  $\Rightarrow$  (ii) Note that  $R/I$  is a non-reduced ring in which any non-zero proper ideal is a prime ideal. Hence we obtain from Lemma 2.6 that  $R/I$  is a quasilocal ring with  $\sqrt{I}/I$  as its unique maximal ideal,  $(\sqrt{I}/I)^2 = I/I$ , and moreover,  $\sqrt{I}/I = R/I(x + I)$  for any  $x \in \sqrt{I} \setminus I$ . Therefore,  $\sqrt{I}$  is a maximal ideal of  $R$ ,  $(\sqrt{I})^2 \subseteq I$ , and  $\sqrt{I} = Rx + I$  for any  $x \in \sqrt{I} \setminus I$ .

(ii)  $\Rightarrow$  (iii) Since  $I \subset \sqrt{I}$ , it follows that  $I$  is not a maximal ideal of  $R$ . Let  $A$  be any proper ideal of  $R$  such that  $I \subset A$ . From  $(\sqrt{I})^2 \subseteq I \subset A$ , it follows that  $\sqrt{I} \subseteq \sqrt{A}$ . Since  $\sqrt{I}$  is a maximal ideal of  $R$ , we obtain  $\sqrt{I} = \sqrt{A}$ . Let  $a \in A \setminus I$ . Then  $a \in \sqrt{I}$ . Hence  $\sqrt{I} = Ra + I \subseteq A$  and so  $A = \sqrt{I}$  is a maximal ideal of  $R$ . This proves that  $I$  is a maximal non-maximal ideal of  $R$ .

(iii)  $\Rightarrow$  (i) As  $I \subset \sqrt{I}$ , it follows that  $I$  is not a prime ideal of  $R$ . Let  $A$  be any proper ideal of  $R$  with  $I \subset A$ . Then  $A$  is a maximal ideal and hence is a prime ideal of  $R$ . This shows that  $I$  is a maximal non-prime ideal of  $R$ .  $\square$

We next proceed to characterize proper ideals  $I$  of a ring  $R$  such that  $I \neq \sqrt{I}$  and  $I$  is a maximal non-primary ideal of  $R$ .

**Proposition 4.2.** *Let  $I$  be a proper ideal of a ring  $R$  such that  $I \neq \sqrt{I}$ . Then the following statements are equivalent:*

- (i)  $I$  is a maximal non-primary ideal of  $R$ .
- (ii)  $\sqrt{I}$  is a prime ideal of  $R$ ,  $R/I$  is quasilocal,  $\dim(R/I) = 1$ , and  $\sqrt{I}/I$  is a simple  $R/I$ -module.

*Proof.* (i)  $\Rightarrow$  (ii) As  $I \neq \sqrt{I}$  and  $I$  is a maximal non-primary ideal of  $R$ , it follows that  $I$  is not a primary ideal of  $R$ , whereas  $\sqrt{I}$  is a primary ideal of  $R$ . Hence  $\sqrt{\sqrt{I}} = \sqrt{I}$  is a prime ideal of  $R$ . Let us denote  $\sqrt{I}$  by  $P$ . Note that  $R/I$  is not a reduced ring, the zero-ideal of  $R/I$  is not primary but each proper nonzero ideal of  $R/I$  is primary. Hence we obtain from Lemma 2.3 that  $R/I$  is quasilocal,  $\dim(R/I) \leq 1$ , and moreover, it follows from Lemma 2.4 that  $P/I$  is a minimal ideal of  $R/I$  (that is,  $P/I$  is a simple  $R/I$ -module). Let  $M/I$  denote the unique maximal ideal of  $R/I$ . Since  $I$  is not a primary ideal of  $R$ , it follows from [1, Proposition 4.2] that  $\sqrt{I}$  is not a maximal ideal of  $R$ . Therefore,  $P/I \subset M/I$  and so  $\dim(R/I) = 1$ .

(ii)  $\Rightarrow$  (i) Note that the ring  $R/I$  satisfies the hypotheses of Lemma 2.5. Hence it follows from Lemma 2.5 that the zero-ideal of  $R/I$  is not a primary ideal. Hence  $I$  is not a primary ideal of  $R$ . Let  $A$  be any proper ideal of  $R$  such that  $I \subset A$ . We consider two cases:

**Case(1)**  $A \subseteq \sqrt{I}$

In this case  $A/I$  is a nonzero ideal of  $R/I$  and  $A/I \subseteq \sqrt{I}/I$ . As  $\sqrt{I}/I$  is a minimal ideal of  $R/I$ , we obtain that  $A/I = \sqrt{I}/I$  and so  $A = \sqrt{I}$  is a prime ideal of  $R$ . Hence  $A$  is a primary ideal of  $R$ .

**Case(2)**  $A \not\subseteq \sqrt{I}$

Let us denote the unique maximal ideal of  $R/I$  by  $M/I$ . Note that  $M$  is the only prime ideal of  $R$  containing  $A$ . Hence it follows that  $\sqrt{A} = M$ . Since  $M$  is a maximal ideal of  $R$ , we obtain from [1, Proposition 4.2] that  $A$  is a primary ideal of  $R$ .

This proves that  $I$  is a maximal non-primary ideal of  $R$ .  $\square$

Recall from [1, p.52] that a proper ideal  $I$  of a ring  $R$  is said to be decomposable if  $I$  admits a primary decomposition (that is,  $I$  can be expressed as the intersection of a finite number of primary ideals of  $R$ ). The following proposition characterizes decomposable ideals  $I$  of a ring  $R$  such that  $I \neq \sqrt{I}$  and  $I$  is a maximal non-primary ideal.

**Proposition 4.3.** *Let  $I$  be a proper ideal of a ring  $R$  such that  $I \neq \sqrt{I}$  and  $I$  is decomposable. The following statements are equivalent:*

- (i)  $I$  is a maximal non-primary ideal of  $R$ .
- (ii)  $\sqrt{I}$  is a prime ideal of  $R$ ,  $(R/I, M/I)$  is quasilocal,  $\dim(R/I) = 1$ ,  $I = \sqrt{I} \cap q$ , where  $q$  is a  $M$ -primary ideal of  $R$ ,  $q \neq M$ , and  $\sqrt{I}/I$  is a simple  $R/I$ -module.

*Proof.* (i)  $\Rightarrow$  (ii) It follows from (i)  $\Rightarrow$  (ii) of Proposition 4.2 that  $\sqrt{I}$  is a prime ideal of  $R$ ,  $R/I$  is quasilocal,  $\dim(R/I) = 1$ , and  $\sqrt{I}/I$  is a simple  $R/I$ -module. Let  $M/I$  denote the unique maximal ideal of  $R/I$ .



We are assuming that  $I$  is decomposable. Let  $I = q_1 \cap \cdots \cap q_n$  be an irredundant primary decomposition of  $I$  in  $R$  with  $q_i$  is a  $P_i$ -primary ideal of  $R$  for each  $i \in \{1, \dots, n\}$ . Since  $I$  is not a primary ideal of  $R$ , it follows that  $n \geq 2$ . Note that  $\sqrt{I} = \bigcap_{i=1}^n P_i$ . As  $\sqrt{I}$  is a prime ideal of  $R$ , it follows that  $\sqrt{I} = P_i$  for some  $i \in \{1, 2, \dots, n\}$ . Without loss of generality we may assume that  $\sqrt{I} = P_1$ . Since  $P_i \neq P_j$  for all distinct  $i, j \in \{1, 2, \dots, n\}$ , it follows that  $P_1 \subset P_j$  for all  $j \in \{2, \dots, n\}$ . As  $P_1/I$  and  $M/I$  are the only prime ideals of  $R/I$ , it follows that  $n = 2$  and  $P_2 = M$ . Note that  $I \subseteq P_1 \cap q_2$ . We assert that  $I = P_1 \cap q_2$ . Since  $q_1 \not\subseteq q_2$ , it follows that  $P_1 \not\subseteq q_2$ . Let  $x \in P_1 \setminus q_2$  and let  $y \in q_2 \setminus P_1$ . Observe that  $xy \in P_1 \cap q_2$  but no power of  $y$  belongs to  $P_1 \cap q_2$  and  $x \notin P_1 \cap q_2$ . Hence  $P_1 \cap q_2$  is not a primary ideal of  $R$ . As we are assuming that  $I$  is a maximal non-primary ideal of  $R$ , it follows that  $I = P_1 \cap q_2$ . Since  $I \neq \sqrt{I}$ , it follows that  $q_2 \neq M$ .  
(ii)  $\Rightarrow$  (i) This follows immediately from (ii)  $\Rightarrow$  (i) of Proposition 4.2.  $\square$

**Example 4.4.** Let  $R = K[[X, Y]]$  be the power series ring in two variables  $X, Y$  over a field  $K$ . It is well-known that  $R$  is a local ring with  $M = RX + RY$  as its unique maximal ideal. Let  $I = RX^2 + RXY$ . Observe that  $I = RX \cap M^2$ . Note that  $\sqrt{I} = RX$  is a prime ideal of  $R$ ,  $M^2 \neq M$  is a  $M$ -primary ideal of  $R$ ,  $\dim(R/I) = 1$ , and  $RX/I$  is a simple  $R/I$ -module. Hence it follows from (ii)  $\Rightarrow$  (i) of Proposition 4.3 that  $I$  is a maximal non-primary ideal of  $R$ .

## 5. MAXIMAL NON-IRREDUCIBLE IDEALS

Recall that an ideal  $I$  of a ring  $R$  is irreducible, if  $I$  is not the intersection of any ideals  $I_1, I_2$  of  $R$  with  $I \subset I_i$  for each  $i \in \{1, 2\}$ . The aim of this section is to determine proper ideals  $I$  of a ring  $R$  such that  $I$  is a maximal non-irreducible ideal of  $R$ . We first characterize proper radical ideals  $I$  of  $R$  such that  $I$  is a maximal non-irreducible ideal of  $R$ .

**Proposition 5.1.** *Let  $I$  be a proper radical ideal of a ring  $R$ . Then the following statements are equivalent:*

- (i)  $I$  is a maximal non-irreducible ideal of  $R$ .
- (ii)  $I = M_1 \cap M_2$  for some distinct maximal ideals  $M_1, M_2$  of  $R$ .

*Proof.* (i)  $\Rightarrow$  (ii) Since  $I$  is a proper radical ideal of  $R$ , it follows from [1, Proposition 1.14] that  $I$  is the intersection of all the prime ideals  $P$  of  $R$  such that  $P \supseteq I$ . Let  $C$  be the collection of all prime ideals  $P$  of  $R$  such that  $P$  is minimal over  $I$ . Observe that we obtain from [5, Theorem 10] that  $I$  is the intersection of all members of  $C$ . Since  $I$  is not irreducible



and any prime ideal is irreducible, we get that  $C$  contains at least two elements. Let  $P_1, P_2 \in C$  be distinct. We assert that  $C = \{P_1, P_2\}$ . Suppose that there exists  $P_3 \in C$  such that  $P_3 \notin \{P_1, P_2\}$ . Then it is clear that  $I \subset P_2 \cap P_3$  and  $P_2 \cap P_3$  is non-irreducible. This is in contradiction to the assumption that  $I$  is a maximal non-irreducible ideal of  $R$ . Therefore,  $C = \{P_1, P_2\}$  and so  $I = P_1 \cap P_2$ . We next show that  $P_1$  and  $P_2$  are maximal ideals of  $R$ . Towards showing it, we first prove that  $P_1 + P_2 = R$ . Suppose that  $P_1 + P_2 \neq R$ . Let  $M$  be a maximal ideal of  $R$  such that  $P_1 + P_2 \subseteq M$ . Since  $P_1$  and  $P_2$  are not comparable under the inclusion relation, there exist  $a \in P_1 \setminus P_2$  and  $b \in P_2 \setminus P_1$ . Consider the ideals  $J_1 = I + Ra + Rb^2$  and  $J_2 = I + Ra^2 + Rb$  of  $R$ . It is clear that  $I \subseteq J_1 \cap J_2$ . As  $a^2 \in (J_1 \cap J_2) \setminus I$ , it follows that  $I \subset J_1 \cap J_2$ . Since  $I$  is a maximal non-irreducible ideal of  $R$ , we obtain that  $J_1 \cap J_2$  is irreducible. Therefore, either  $J_1 \subseteq J_2$  or  $J_2 \subseteq J_1$ . If  $J_1 \subseteq J_2$ , then  $a = x + ra^2 + sb$  for some  $x \in I = P_1 \cap P_2$  and  $r, s \in R$ . This implies that  $a(1 - ra) = x + sb \in P_2$ . As  $a \notin P_2$ , we obtain that  $1 - ra \in P_2$ . Therefore,  $1 = ra + 1 - ra \in P_1 + P_2 \subseteq M$ . This is a contradiction. Observe that we get a similar contradiction if  $J_2 \subseteq J_1$ . Hence  $P_1 + P_2 = R$ . Let  $M_1$  be a maximal ideal of  $R$  such that  $P_1 \subseteq M_1$ . Since  $P_1 + P_2 = R$ , it follows that the ideal  $M_1 \cap P_2$  is not irreducible. As  $I \subseteq M_1 \cap P_2$ , we obtain that  $I = P_1 \cap P_2 = M_1 \cap P_2$ . Since  $P_1 \not\subseteq P_2$ , it follows that  $P_1 \supseteq M_1$  and so  $P_1 = M_1$  is a maximal ideal of  $R$ . Similarly it can be shown that  $P_2$  is a maximal ideal of  $R$ . Thus  $I = M_1 \cap M_2$  for some distinct maximal ideals  $M_1, M_2$  of  $R$ .

(ii)  $\Rightarrow$  (i) If  $I = M_1 \cap M_2$  for some distinct maximal ideals  $M_1, M_2$  of  $R$ , then it is clear that  $I$  is not irreducible. It is verified in the proof of (ii)  $\Rightarrow$  (iii) of Proposition 3.2 that  $M_1$  and  $M_2$  are the only proper ideals  $J$  of  $R$  such that  $I \subset J$ . Since  $M_1$  and  $M_2$  are both irreducible, we obtain that  $I$  is a maximal non-irreducible ideal of  $R$ .  $\square$

Let  $I$  be a proper ideal of a ring  $R$  such that  $I \neq \sqrt{I}$ . We next attempt to characterize such ideals  $I$  in order that  $I$  is a maximal non-irreducible ideal of  $R$ . We do not know the precise characterization of such ideals. However, we have the following partial results.

**Lemma 5.2.** *Let  $I$  be a proper ideal of a ring  $R$  such that  $I \neq \sqrt{I}$ . If  $I$  is a maximal non-irreducible ideal of  $R$ , then  $\sqrt{I}$  is a prime ideal of  $R$  and moreover,  $R/I$  is quasilocal.*

*Proof.* Let  $C$  be the collection of all prime ideals  $P$  of  $R$  such that  $P$  is minimal over  $I$ . We assert that  $C$  is singleton. Let  $P, Q \in C$ . Since  $I \neq \sqrt{I}$ , it is clear that  $I \subset P \cap Q$ . As  $I$  is a maximal non-irreducible ideal of  $R$ , it follows that  $P \cap Q$  is irreducible. Hence either  $P \subseteq Q$  or

$Q \subseteq P$ . Therefore,  $P = Q$ . This shows that there is only one prime ideal  $P$  of  $R$  such that  $P$  is minimal over  $I$ . Thus  $\sqrt{I} = P$  is a prime ideal of  $R$ .

We next show that  $R/I$  is quasilocal. Let  $M, N$  be maximal ideals of  $R$  such that  $I \subseteq M \cap N$ . Since  $I \neq \sqrt{I}$ , it follows that  $I \subset M \cap N$ . As  $M \cap N$  is irreducible, we obtain that either  $M \subseteq N$  or  $N \subseteq M$ . Hence  $M = N$ . This shows that  $R/I$  is quasilocal.  $\square$

**Lemma 5.3.** *Let  $(T, N)$  be a quasilocal ring such that  $(0) \neq \sqrt{(0)}$  and  $(0)$  is a maximal non-irreducible ideal of  $T$ . Then  $\dim_{T/N}(N/N^2) \leq 2$ .*

*Proof.* Suppose that  $\dim_{T/N}(N/N^2) \geq 3$ . Let  $\{a, b, c\} \subseteq N$  be such that  $\{a+N^2, b+N^2, c+N^2\}$  is linearly independent over  $T/N$ . Consider the ideals  $J_1 = Ta + Tc$  and  $J_2 = Tb + Tc$ . By the choice of  $a, b, c$ , it is clear that  $J_1 \not\subseteq J_2$ ,  $J_2 \not\subseteq J_1$  and so  $J_1 \cap J_2$  is not an irreducible ideal of  $T$ . Moreover, as  $c \in J_1 \cap J_2$ , it follows that  $J_1 \cap J_2 \neq (0)$ . This contradicts the hypothesis that  $(0)$  is a maximal non-irreducible ideal of  $T$ . Therefore,  $\dim_{T/N}(N/N^2) \leq 2$ .  $\square$

**Lemma 5.4.** *Let  $(T, N)$  be a quasilocal ring such that  $(0) \neq \sqrt{(0)}$  and  $\dim_{T/N}(N/N^2) = 2$ . Then the following statements are equivalent:*

- (i)  $(0)$  is a maximal non-irreducible ideal of  $T$ .
- (ii)  $N^2 = (0)$ .

*Proof.* By hypothesis,  $\dim_{T/N}(N/N^2) = 2$ . Let  $\{a, b\} \subseteq N$  be such that  $\{a + N^2, b + N^2\}$  is a basis of  $N/N^2$  as a vector space over  $T/N$ .  
 (i)  $\Rightarrow$  (ii) Consider the ideals  $J_1 = N^2 + Ta$  and  $J_2 = N^2 + Tb$ . By the choice of the elements  $a, b$ , it is clear that  $J_1 \not\subseteq J_2$  and  $J_2 \not\subseteq J_1$ . Hence the ideal  $J_1 \cap J_2$  is not irreducible. Since  $(0)$  is a maximal non-irreducible ideal of  $T$ , it follows that  $J_1 \cap J_2 = (0)$ . As  $N^2 \subseteq J_1 \cap J_2$ , we obtain that  $N^2 = (0)$ .

(ii)  $\Rightarrow$  (i) It follows from  $N^2 = (0)$  and from the choice of the elements  $a, b$  that  $Ta \not\subseteq Tb$ ,  $Tb \not\subseteq Ta$ , and  $Ta \cap Tb = (0)$ . This implies that  $(0)$  is not an irreducible ideal of  $T$ . Let  $J$  be any nonzero proper ideal of  $T$ . Then either  $\dim_{T/N}(J) = 1$  or  $2$ . If  $\dim_{T/N}(J) = 2$ , then  $J = N$  is irreducible. Suppose that  $\dim_{T/N}(J) = 1$ . Let  $A, B$  be proper ideals of  $T$  such that  $J = A \cap B$ . If  $J \neq A$  and  $J \neq B$ , then we get that  $A = B = N$  and so  $J = N$ . This is a contradiction. Hence either  $J = A$  or  $J = B$ . This shows that  $J$  is irreducible. Hence  $(0)$  is a maximal non-irreducible ideal of  $T$ .  $\square$

## Acknowledgements

We are very much thankful to the referee for a very careful reading of this paper and valuable suggestions. We are also very much thankful to Professor H. Ansari-Toroghy for his support.

## REFERENCES

- [1] M.F. Atiyah and I.G. Macdonald, *Introduction to commutative algebra*. Addison-Wesley, 1969.
- [2] A. Ayache, D. E. Dobbs, and O. Echi, *On maximal non-ACCP subrings*, J. Algebra Appl. (5)**6** (2007), 873-894.
- [3] A. Ayache and N. Jarboui, *Maximal non-Noetherian subring of a domain*, J. Algebra **248** (2002), 806-823.
- [4] A. Ayache, M. Ben Nasr, and N. Jarboui, *PID pairs of rings and maximal non-PID subrings*, Math. Z. **260** (2011), 635-647.
- [5] I. Kaplansky, *Commutative Rings*. The University of Chicago Press, Chicago, 1974.

**S. Visweswaran**

Department of Mathematics, Saurashtra University, P.O. Box No. 360 005 Rajkot, India.

Email: [s\\_visweswaran2006@yahoo.co.in](mailto:s_visweswaran2006@yahoo.co.in)

**Anirudhdha Parmar**

Department of Mathematics, Saurashtra University, P.O. Box No. 360 005 Rajkot, India.

Email: [anirudh.maths@gmail.com](mailto:anirudh.maths@gmail.com)